

On the domination of Cartesian product of directed cycles: Results for certain equivalence classes of lengths

Michel Mollard*

Institut Fourier
100, rue des Maths
38402 St Martin d'Hères Cedex FRANCE
`michel.mollard@ujf-grenoble.fr`

Abstract

Let $\gamma(\vec{C}_m \square \vec{C}_n)$ be the domination number of the Cartesian product of directed cycles \vec{C}_m and \vec{C}_n for $m, n \geq 2$. Shaheen [13] and Liu et al. ([11], [12]) determined the value of $\gamma(\vec{C}_m \square \vec{C}_n)$ when $m \leq 6$ and [12] when both m and $n \equiv 0 \pmod{3}$. In this article we give, in general, the value of $\gamma(\vec{C}_m \square \vec{C}_n)$ when $m \equiv 2 \pmod{3}$ and improve the known lower bounds for most of the remaining cases. We also disprove the conjectured formula for the case $m \equiv 0 \pmod{3}$ appearing in [12].

AMS Classification[2010]:05C69,05C38.

Keywords: Directed graph, Cartesian product, Domination number, Directed cycle.

1 Introduction and definitions

Let $D = (V, A)$ be a finite directed graph (digraph for short) without loops or multiple arcs.

A vertex u *dominates* a vertex v if $u = v$ or $uv \in A$. A set $W \subseteq V$ is a *dominating set* of D if any vertex of V is dominated by at least one vertex of W . The *domination number* of D , denoted by $\gamma(D)$ is the minimum cardinality of a dominating set. The set V is a dominating set thus $\gamma(D)$ is finite. These definitions extend to digraphs the classical domination notion for undirected graphs.

The determination of the domination number of a directed or undirected graph is, in general, a difficult question in graph theory. Furthermore this problem has

*CNRS Université Joseph Fourier

connections with information theory. For example the domination number of hypercubes is linked to error-correcting codes. Among the lot of related works, Haynes et al. ([7], [8]), mention the special cases of the domination of Cartesian products of undirected paths, cycles or more generally graphs([1] to [6], [9], [10]).

For two digraphs $D_1 = (V_1, A_1)$ and $D_2 = (V_2, A_2)$ the *Cartesian product* $D_1 \square D_2$ is the digraph with vertex set $V_1 \times V_2$ and $(x_1, x_2)(y_1, y_2) \in A(D_1 \square D_2)$ if and only if $x_1 y_1 \in A_1$ and $x_2 = y_2$ or $x_2 y_2 \in A_2$ and $x_1 = y_1$. Note that $D_2 \square D_1$ is isomorphic to $D_1 \square D_2$. In [13] Shaheen determined the domination number of $\overrightarrow{C_m} \square \overrightarrow{C_n}$ for $m \leq 6$ and arbitrary n . In two articles [11], [12] Liu et al. considered independently the domination number of the Cartesian product of two directed cycles. They gave also the value of $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n})$ when $m \leq 6$ and when both m and $n \equiv 0 \pmod{3}$ [12]. Furthermore they proposed lower and upper bounds for the general case.

In this paper we are able to give, in general, the value of $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n})$ when $m \equiv 2 \pmod{3}$ and we improve the lower bounds for most of the still unknown cases. We also disprove the conjectured formula appearing in [12] for the case $m \equiv 0 \pmod{3}$.

We denote the vertices of a directed cycle $\overrightarrow{C_n}$ by $C_n = \{0, 1, \dots, n-1\}$, the integers considered modulo n . Thus, when used for vertex labeling, $a+b$ and $a-b$ will denote the vertices $a+b$ and $a-b \pmod{n}$. Notice that there exists an arc xy from x to y in $\overrightarrow{C_n}$ if and only if $y \equiv x+1 \pmod{n}$, thus with our convention, if and only if $y = x+1$. For any i in $\{0, 1, \dots, n-1\}$ we will denote by $\overrightarrow{C_m^i}$ the subgraph of $\overrightarrow{C_m} \square \overrightarrow{C_n}$ induced by the vertices $\{(k, i) \mid k \in \{0, 1, \dots, m-1\}\}$. Note that $\overrightarrow{C_m^i}$ is isomorphic to $\overrightarrow{C_m}$. We will denote by C_m^i the set of vertices of $\overrightarrow{C_m^i}$.

2 General bounds and the case $m \equiv 2 \pmod{3}$

We start this section by developing a general upper bound for $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n})$. Then we will construct minimum dominating sets for $m \equiv 2 \pmod{3}$. These optimal sets will be obtained from integer solutions of a system of equations.

Proposition 1 *Let W be a dominating set of $\overrightarrow{C_m} \square \overrightarrow{C_n}$. Then for all i in $\{0, 1, \dots, n-1\}$ considered modulo n we have $|W \cap C_m^{i-1}| + 2|W \cap C_m^i| \geq m$.*

Proof : The m vertices of C_m^i can only be dominated by vertices of $W \cap C_m^i$ and $W \cap C_m^{i-1}$. Each of the vertices of $W \cap C_m^i$ dominates two vertices in C_m^i . Similarly each of the vertices of $W \cap C_m^{i-1}$ dominates one vertex in C_m^i . The result follows. \square

Theorem 2 *Let $m, n \geq 2$ and $k_1 = \lfloor \frac{m}{3} \rfloor$ then*

- (i) *if $m \equiv 0 \pmod{3}$ then $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) \geq nk_1$, or*
- (ii) *if $m \equiv 1 \pmod{3}$ then $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) \geq nk_1 + \frac{n}{2}$, or*
- (iii) *if $m \equiv 2 \pmod{3}$ then $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) \geq nk_1 + n$.*

Proof :

Let W be a dominating set of $\overrightarrow{C_m} \square \overrightarrow{C_n}$ and for any i in $\{0, 1, \dots, n-1\}$ let $a_i = |W \cap C_m^i|$. Notice first, as noticed by Liu et al.([12]), that each of the vertices of W dominates three vertices of $\overrightarrow{C_m} \square \overrightarrow{C_n}$ and thus $|W| \geq \frac{mn}{3}$. This general bound give the announced result for $m = 3k_1$, $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) \geq nk_1 + \frac{n}{3}$ for $m = 3k_1 + 1$ and $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) \geq nk_1 + 2\frac{n}{3}$ for $m = 3k_1 + 2$. We will improve these two last results to verify parts (ii) and (iii) of the theorem.

Assume first $m = 3k_1 + 1$. Let J be the set of $j \in \{0, 1, \dots, n-1\}$ such that $a_j \leq k_1$. If $J = \emptyset$ then $|W| \geq n(k_1 + 1) \geq nk_1 + \frac{n}{2}$ and we are done. Otherwise let $J' = \{j \mid j+1 \pmod n \in J\}$. By Proposition 1, for any i in $\{0, 1, \dots, n-1\}$ considered modulo n , we have $a_{i-1} + 2a_i \geq 3k_1 + 1$. Then if i belongs to J , $a_{i-1} + a_i \geq 2k_1 + 1$. A first consequence is that there are no consecutive indices, taken modulo n , in J . Indeed if $j-1$ and j are in J then, by definition of J , $a_{j-1} + a_j \leq 2k_1$ in contradiction with the previous inequality. By definition of J' we have thus $J \cap J' = \emptyset$.

Now let $K = \{j \in \{0, 1, \dots, n-1\} \mid j \notin J \cup J'\}$. We can write $\{0, 1, \dots, n-1\} = J \cup J' \cup K$ where J, J' and K are disjoint sets. Notice that $\theta : j \mapsto j-1 \pmod n$ induces a one to one mapping between J and J' .

The cardinality of W is $|W| = \sum_{i \in \{0, 1, \dots, n-1\}} a_i = \sum_{i \in J} a_i + \sum_{i \in J'} a_i + \sum_{i \in K} a_i$. We can use θ for grouping 2 by 2 the elements of $J \cup J'$ and write $\sum_{i \in J} a_i + \sum_{i \in J'} a_i = \sum_{i \in J} a_i + \sum_{i \in J} a_{\theta(i)} = \sum_{i \in J} (a_i + a_{i-1})$. Using $a_{i-1} + a_i \geq 2k_1 + 1$, because $i \in J$, we obtain $\sum_{i \in J} a_i + \sum_{i \in J'} a_i \geq |J|(2k_1 + 1)$.

If $i \in K$ then $i \notin J$ and $a_i \geq k_1 + 1$. Since $|K| = n - 2|J|$ we have $\sum_{i \in K} a_i \geq (n - 2|J|)(k_1 + 1)$. Then $|W| = \sum_{i \in \{0, 1, \dots, n-1\}} a_i \geq |J|(2k_1 + 1) + (n - 2|J|)(k_1 + 1) = nk_1 + n - |J|$.

Since $|J| = |J'|$ and $J \cap J' = \emptyset$, $n - |J| \geq \frac{n}{2}$ and the conclusion for (ii) follows.

The case $m = 3k_1 + 2$ is similar. Let J be the set of $j \in \{0, 1, \dots, n-1\}$ such that $a_j \leq k_1$. If $J = \emptyset$ then we are done. Otherwise let $J' = \{j \mid j+1 \pmod n \in J\}$. If $i \in J$ we have $a_{i-1} + 2a_i \geq 3k_1 + 2$ thus $a_{i-1} + a_i \geq 2k_1 + 2$. Then $J \cap J' = \emptyset$ and $\sum_{i \in J \cup J'} a_i \geq |J|(2k_1 + 2)$. Therefore $\sum_{i \in \{0, 1, \dots, n-1\}} a_i \geq |J|(2k_1 + 2) + (n - 2|J|)(k_1 + 1) \geq n(k_1 + 1)$.

□

Let us now study in detail the case $m \equiv 2 \pmod 3$. Assume $m = 3k_1 + 2$. Let A be the set of $k_1 + 1$ vertices of $\overrightarrow{C_m}$ defined by $A = \{0\} \cup \{2 + 3p \mid p = 0, 1, \dots, k_1 - 1\} = \{0\} \cup \{2, 5, \dots, m-6, m-3\}$. For any i in $\{0, 1, \dots, m-1\}$ let us call $A_i = \{j \mid j-i \pmod m \in A\}$ the *translate*, considered modulo m , of A by i . We have thus $A_i = \{i\} \cup \{i+2, i+5, \dots, i-6, i-3\}$ (see Figure 1).

We will call a set S of vertices of $\overrightarrow{C_m} \square \overrightarrow{C_n}$ a A -set if for any j in $\{0, 1, \dots, n-1\}$ we have $S \cap C_m^j = A_i$ for some i in $\{0, 1, \dots, n-1\}$. It will be convenient to denote this index i , function of j , as i_j . If S is a A -set then $|S| = n(k_1 + 1)$; thus if a set is both a A -set and a dominating set, by Theorem 2, it is minimum and we have $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) = n(k_1 + 1)$.

Lemma 3 *Let $m = 3k_1 + 2$. Let S be a A -set and for any j in $\{0, 1, \dots, n-1\}$*




Figure 1: A_i, A_{i-1} and A_{i+2}

define i_j as the index such that $S \cap C_m^j = A_{i_j}$. Assume that:

$$\begin{cases} \text{for any } j \in \{1, \dots, n-1\} & i_j \equiv i_{j-1} + 1 \pmod{m} \text{ or } i_j \equiv i_{j-1} - 2 \pmod{m} \\ \text{and} \\ i_0 \equiv i_{n-1} + 1 \pmod{m} \text{ or } i_0 \equiv i_{n-1} - 2 \pmod{m}. \end{cases}$$

then S is a dominating set of $\overrightarrow{C_m} \square \overrightarrow{C_n}$.

Proof :

Note first that for any i in $\{0, 1, \dots, m-1\}$ the set of non dominated vertices of C_m by A_i is $T = \{i+4, i+7, \dots, i-4, i-1\}$. Note also that $A_{i+2} = \{i+2\} \cup \{i+4, i+7, \dots, i-4, i-1\}$ and $A_{i-1} = \{i-1\} \cup \{i+1, i+4, \dots, i-7, i-4\}$. Thus $T \subset A_{i+2}$ and $T \subset A_{i-1}$.

Let j in $\{1, \dots, n-1\}$. Let us prove that the vertices of C_m^j are dominated. Indeed, by the previous remark and the lemma hypothesis, the vertices non dominated by $S \cap C_m^j$ are dominated by $S \cap C_m^{j-1}$ (see Figure 1). For the same reasons the vertices of C_m^0 are dominated by those of $S \cap C_m^0$ and $S \cap C_m^{n-1}$.

□

We will prove next that the existence of solutions to some system of equations over integers implies the existence of a A -set satisfying the hypothesis of Lemma 3.

Lemma 4 Let $m = 3k_1 + 2$. If there exist integers $a, b \geq 0$ such that

$$\begin{cases} a + b = n - 1 \\ \text{and} \\ a - 2b \equiv 2 \pmod{m} \text{ or } a - 2b \equiv m - 1 \pmod{m} \end{cases}$$

then $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n}) = n(k_1 + 1)$.

Proof : Consider a word $w = w_1 \dots w_{n-1}$ on the alphabet $\{1, -2\}$ with a occurrences of 1 and b of -2 . Such a word exists, for example $w = 1^a(-2)^b$. We can associate with w a set S of vertices of $\vec{C}_m \square \vec{C}_n$ using the following algorithm:

- $S \cap C_m^0 = A_0$
- For $i = 1$ to $n - 1$ do
 - begin
 - Let k such that $S \cap C_m^{i-1} = A_k$
 - If $w_i = 1$ let $k' \equiv k + 1 \pmod{m}$ else $k' \equiv k - 2 \pmod{m}$
 - $S \cap C_m^i := A_{k'}$
 - end

By construction S is a A -set. Notice that we have $S \cap C_m^{n-1} := A_{i_{n-1}}$ where $i_{n-1} \equiv \sum_{k=1}^{n-1} w_k \equiv a - 2b \pmod{m}$. Thus $i_{n-1} \equiv 2 \pmod{m}$ or $i_{n-1} \equiv m - 1 \pmod{m}$. By Lemma 3 S is a dominating set. Furthermore, because S is a A -set, $|S| = n(k_1 + 1)$, thus by Theorem 2 it is minimum and we have $\gamma(\vec{C}_m \square \vec{C}_n) = n(k_1 + 1)$. \square
 With the exception of one sub case we can find solutions (a, b) of the system and thus obtain minimum dominating sets for $m \equiv 2 \pmod{3}$.

Theorem 5 Let $m, n \geq 2$ and $m \equiv 2 \pmod{3}$. Let $k_1 = \lfloor \frac{m}{3} \rfloor$ and $k_2 = \lfloor \frac{n}{3} \rfloor$.

- (i) if $n = 3k_2$ then $\gamma(\vec{C}_m \square \vec{C}_n) = n(k_1 + 1)$, and
- (ii) if $n = 3k_2 + 1$ and $2k_2 \geq k_1$ then $\gamma(\vec{C}_m \square \vec{C}_n) = n(k_1 + 1)$, and
- (iii) if $n = 3k_2 + 1$ and $2k_2 < k_1$ then $\gamma(\vec{C}_m \square \vec{C}_n) > n(k_1 + 1)$, and
- (iv) if $n = 3k_2 + 2$ and $n \geq m$ then $\gamma(\vec{C}_m \square \vec{C}_n) = n(k_1 + 1)$, and
- (v) if $n = 3k_2 + 2$ and $n \leq m$ then $\gamma(\vec{C}_m \square \vec{C}_n) = m(k_2 + 1)$.

Proof : We will use Lemma 4 considering the following integer solutions of

$$\begin{cases} a, b \geq 0 \\ a + b = n - 1 \\ a - 2b \equiv 2 \pmod{m} \text{ or } a - 2b \equiv m - 1 \pmod{m}. \end{cases}$$

- (i) if $n = 3k_2$ then $k_2 \geq 1$. Take $a = 2k_2$ and $b = k_2 - 1$.
- (ii) if $n = 3k_2 + 1$ and $2k_2 \geq k_1$ then take $a = 2k_2 - k_1$ and $b = k_2 + k_1$.
- (iii) if $n = 3k_2 + 1$ and $2k_2 < k_1$ then $\gamma(\vec{C}_m \square \vec{C}_n) = \gamma(\vec{C}_n \square \vec{C}_m) \geq \frac{(2k_2+1)m}{2}$ by Theorem 2. Furthermore $\frac{(2k_2+1)m}{2} - n(k_1 + 1) = \frac{k_1}{2} - k_2 > 0$.
- (iv) if $n = 3k_2 + 2$ and $k_2 \geq k_1$ then take $a = 2k_2 - 2k_1$ and $b = k_2 + 2k_1 + 1$.
- (v) if $n = 3k_2 + 2$ and $k_2 \leq k_1$ then use $\gamma(\vec{C}_m \square \vec{C}_n) = \gamma(\vec{C}_n \square \vec{C}_m)$.

\square

3 The case $m \equiv 0 \pmod{3}$

In [12] Liu et al. conjectured the following formula:

Conjecture 6 *Let $k \geq 2$. Then $\gamma(\overrightarrow{C_{3k}} \square \overrightarrow{C_n}) = k(n+1)$ for $n \not\equiv 0 \pmod{3}$.*

Our Theorem 5 confirms the conjecture when $n \equiv 2 \pmod{3}$. Unfortunately the formula is not always valid when $n \equiv 1 \pmod{3}$.

Indeed consider $C_{3k} \square C_4$. In [11] the following result is proved:

Theorem 7 *Let $n \geq 2$. Then $\gamma(\overrightarrow{C_4} \square \overrightarrow{C_n}) = \frac{3n}{2}$ if $n \equiv 0 \pmod{8}$ and $\gamma(\overrightarrow{C_4} \square \overrightarrow{C_n}) = n + \lceil \frac{n+1}{2} \rceil$ otherwise.*

We have thus $\gamma(\overrightarrow{C_{3k}} \square \overrightarrow{C_4}) = \gamma(\overrightarrow{C_4} \square \overrightarrow{C_{3k}}) = 3k + \lceil \frac{3k+1}{2} \rceil$ when $k \not\equiv 0 \pmod{8}$. Alternately, Conjecture 6 proposes the value $\gamma(\overrightarrow{C_{3k}} \square \overrightarrow{C_4}) = 5k$. These two numbers are different when $k \geq 3$.

4 Conclusion

Consider the possible remainder of m, n modulo 3. For some of the nine possibilities, we have found exact values for $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n})$. The remaining cases are:

- a) $m \equiv 0 \pmod{3}$ and $n \equiv 1 \pmod{3}$
- b) The symmetrical case $m \equiv 1 \pmod{3}$ and $n \equiv 0 \pmod{3}$.
- c) m and $n \equiv 1 \pmod{3}$.
- d) The case m or $n \equiv 2 \pmod{3}$ is not completely solved by Theorem 5. The following subcases are still open
 - i) $m \equiv 2 \pmod{3}$ and $n \equiv 1 \pmod{3}$ with $m > 2n + 1$
 - ii) the symmetrical case $m \equiv 1 \pmod{3}$ and $n \equiv 2 \pmod{3}$ with $n > 2m + 1$.

For these values of m, n there does not always exist a dominating set reaching the bound stated in Theorem 2 and thus the determination of $\gamma(\overrightarrow{C_m} \square \overrightarrow{C_n})$ seems to be a more difficult problem.

References

- [1] T.Y. Chang, W.E. Clark: "The Domination numbers of the $5 \times n$ and $6 \times n$ grid graphs", *J. Graph Theory*, **17** (1993) 81-107.
- [2] M. El-Zahar, C.M. Pareek: "Domination number of products of graphs", *Ars Combin.*, **31** (1991) 223-227.
- [3] M. El-Zahar, S. Khamis, Kh. Nazzal: "On the Domination number of the Cartesian product of the cycle of length n and any graph", *Discrete App. Math.*, **155** (2007) 515-522.

- [4] R.J. Faudree, R.H. Schelp: ‘The Domination number for the product of graphs’, *Congr. Numer.*, **79** (1990) 29-33.
- [5] S. Gravier, M. Mollard: “On Domination numbers of Cartesian product of paths”, *Discrete App. Math.*, **80** (1997) 247-250.
- [6] B. Hartnell, D. Rall: “On dominating the Cartesian product of a graph and K_2 ”, *Discuss. Math. Graph Theory*, **24(3)** (2004) 389-402.
- [7] T.W. Haynes, S.T. Hedetniemi, and P.J. Slater: *Fundamentals of Domination in Graphs*, Marcel Dekker, Inc. New York, 1998.
- [8] T.W. Haynes, S.T. Hedetniemi, and P.J. Slater eds.: *Domination in Graphs: Advanced Topics*, Marcel Dekker, Inc. New York, 1998.
- [9] M.S. Jacobson, L.F. Kinch: “On the Domination number of products of graphs I”, *Ars Combin.*, **18** (1983) 33-44.
- [10] S. Klavžar, N. Seifter: “Dominating Cartesian products of cycles”, *Discrete App. Math.*, **59** (1995) 129-136.
- [11] J. Liu, X.D. Zhang, X. Chen, J. Meng: “On domination number of Cartesian product of directed cycles”, *Inf. Process. Lett.*, **110(5)** (2010) 171-173.
- [12] J. Liu, X.D. Zhang, X. Chen, J. Meng: “Domination number of Cartesian products of directed cycles”, *Inf. Process. Lett.*, **111(1)** (2010) 36-39.
- [13] R.S. Shaheen: “Domination number of toroidal grid digraphs”, *Utilitas Mathematica* **78**(2009) 175-184.